



ST. LUKE'S[®] Episcopal Hospital Blood Center

In The Past 48 Hours Have You Taken Aspirin Or Anything That Contains Aspirin?

Aspirin irreversibly inactivates platelet function. Donors who are taking aspirin or any medication containing aspirin should not be the sole source of platelets for a single transfusion dose.

If you have taken any medication that contains aspirin in the last 48 hours:

- **Whole blood donation:** Acceptable to donate whole blood. The platelet bag is marked to indicate aspirin (ASA).
- **Apheresis Platelet Donation:** Not acceptable for platelet donation. Please donate either whole blood or plasma at this time.

THE FOLLOWING COMMONLY USED MEDICATIONS CONTAIN ASPIRIN OR MEDICATIONS KNOWN TO INHIBIT PLATELET FUNCTION:

| | | | |
|--------------------|-------------------------|--------------|------------------|
| 4-Way Cold Tablets | Dipyridamole | Midol | Sine-off |
| Alka-Selzer | Doan's Pill | Norgesic | Soma compound |
| Aspirin | Dristan | Pepto-Bismol | Standback powder |
| Anacin | Ecotrin | Percodan | Ticlid* |
| Baby Aspirin | Excederin | Persantine | Triaminic |
| Bayer | Feldene | Piroxicam | |
| BC Powder | Fiorinal | Plavix* | |
| Bufferin | Goody's Headache Powder | Robaxin | |

* Donors that give a history of taking Ticlid or Plavix should be deferred from donating platelets for 14 days after completion of medication.

THE FOLLOWING MEDICATIONS DO NOT CONTAIN ASPIRIN:

| | | | |
|--------------------|----------------|------------|--------------------|
| Acetaminophen | Dramamine | Nuprin | Tylenol |
| Advil | IB Ibuprofen | Nytol | Tylenol with sinus |
| Aleve | Ibuprofen | Ornex | Tylenol PM |
| Allerest | Kaopectate | Pamprin | Tums |
| Bayer IB Ibuprofen | Malos | Robitussin | Vioxx |
| Celebrex | Motrin | Rolaids | |
| Contact | Naprosyn | Rufen | |
| Darvocet | Neo-Synephrine | Sinutab | |
| Di-Gel | No Doz | Sleep-eze | |
| Dimetapp | | Sudafed | |